

AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

LISTING OF CLAIMS:

1. (Currently amended) An isolated nucleic acid molecule comprising ~~SEQ ID NO:3~~ a nucleotide sequence encoding a haemopoietin receptor comprising ~~[[an]]the~~ amino acid sequence set forth in SEQ ID NO:4 or a nucleotide sequence encoding a derivative of said haemopoietin receptor, wherein the derivative binds with IL-13 or is immunologically interactive with antibodies to an IL-13 receptor alpha chain.

2. (Currently amended) An isolated nucleic acid molecule comprising ~~SEQ ID NO:3~~ a nucleotide sequence encoding a haemopoietin receptor comprising an amino acid sequence as set forth in SEQ ID NO:4 or a nucleotide sequence encoding a derivative of said haemopoietin receptor thereof, wherein said receptor:

(i) binds with IL-13 or its derivatives; and

(ii) binds with a complex between IL-4 and IL-4 receptor α -chain,

and wherein the derivative of said haemopoietin receptor binds with IL-13 or is immunologically interactive with antibodies to an IL-13 receptor alpha chain.

3-6. (Cancelled)

7. (Currently amended) An isolated nucleic acid molecule comprising a sequence of nucleotides which encodes an IL-13 receptor α -chain or a derivative thereof, said nucleic acid molecule having a nucleotide sequence as set forth in SEQ ID NO:3 or having a nucleic acid molecule which hybridizes to the nucleotide sequence as set forth in SEQ ID NO:3 under low stringency conditions, wherein said low stringency conditions comprise 6x SSC, 0.1% w/v SDS at 42°C, and wherein the derivative binds with IL-13 or is immunologically interactive with antibodies to an IL-13 receptor alpha chain.

8. (Currently amended) An isolated nucleic acid molecule comprising a sequence of nucleotides which encodes an IL-13 receptor α -chain or a derivative thereof having an amino acid sequence as set forth in SEQ ID NO:4, wherein the derivative binds with IL-13 or is immunologically interactive with antibodies to an IL-13 receptor alpha chain.

9. (Original) An isolated nucleic acid molecule according to claim 1 or 2 or 7 or 8 which encodes a haemopoietin receptor capable of interaction with IL-13 or its derivatives, which interaction is capable of competitive inhibition by IL-4 or a derivative thereof in cells which express an IL-4 receptor α -chain.

10. (Previously presented) An expression vector comprising a nucleic acid molecule according to claim 1 or 7 operably linked to a promoter which directs expression of said nucleic acid molecule in a host cell.

11-24. (Cancelled)

25. (Previously presented) A composition comprising a nucleic acid molecule according to claim 1 or 2 or 7 or 8 and a pharmaceutically acceptable carrier.

26-27. (Cancelled)

28. (Previously presented) A method of producing a recombinant polypeptide having at least two of the following characteristics:

- (i) comprises an amino acid sequence as set forth in SEQ ID NO:4;
- (ii) is encoded by a nucleotide sequence as set forth in SEQ ID NO:3;
- (iii) binds with IL-13 or its derivatives; and
- (iv) said polypeptide, when expressed in COS cells, has a molecular weight of from about 50,000 to about 70,000 daltons as determined by Western blot analysis,

said method comprising culturing cells comprising the expression vector according to claim 10 for a time and under conditions sufficient to express the nucleic acid molecule in said expression vector to produce a recombinant polypeptide and isolating said recombinant polypeptide.

29. (Previously presented) A method of producing a recombinant polypeptide having at least three of the following characteristics:

- (i) comprises an amino acid sequence as set forth in SEQ ID NO:4;
- (ii) is encoded by a nucleotide sequence as set forth in SEQ ID NO:3;
- (iii) binds with IL-13 or its derivatives;
- (iv) said polypeptide, when expressed in COS cells, has a molecular weight of from about 50,000 to about 70,000 daltons as determined by Western blot analysis;
- (v) comprises an amino acid sequence derived from IL-4 receptor α -chain; and
- (vi) is capable of interaction with IL-13 which is competitively inhibited by IL-4 in cells which express an IL-4 receptor α -chain,

said method comprising culturing cells comprising the expression vector according to claim 10 for a time and under conditions sufficient to express the nucleic acid molecule in said expression vector to produce a recombinant polypeptide and isolating said recombinant polypeptide.

30. (Previously presented) An isolated host cell which expresses the recombinant polypeptide produced by the method according to claim 28.

31-35. (Cancelled)

36. (Previously presented) An isolated host cell which expresses the recombinant polypeptide produced by the method according to claim 29.

37. (Currently amended) An isolated nucleic acid molecule comprising the nucleotide sequence as set forth in SEQ ID NO: 3.

38. (Currently amended) An isolated nucleic acid molecule comprising a sequence of nucleotides which encodes an extracellular domain of an IL-13 ~~haemopoietin~~ receptor alpha chain.

39. (Currently amended) The isolated nucleic acid molecule of claim 38 wherein said ~~extra-cellular~~ extracellular domain is an immunoglobulin-like domain.

40. (Currently amended) The isolated nucleic acid molecule of claim 38 wherein said ~~extra~~ cellular extracellular domain is an haemopoietin receptor domain.
41. (Currently amended) The isolated nucleic acid molecule of claim 39 wherein said immunoglobulin-like domain consists ~~essentially~~ of amino acids 28-118 of SEQ ID NO:4.
42. (Currently amended) The isolated nucleic acid molecule of claim 40 wherein said haemopoietin receptor domain consists ~~essentially~~ of amino acids 119-342 of SEQ ID NO:4
43. (Currently amended) The isolated nucleic acid molecule of claim 1 ~~Claim 37~~, encoding a polypeptide consisting ~~essentially~~ of amino acids 28-346 of SEQ ID NO: 4.
44. (Currently amended) The isolated nucleic acid molecule of claim 1 ~~Claim 37~~, encoding a polypeptide consisting ~~essentially~~ of amino acids 28-426 of SEQ ID NO:4.
45. (Currently amended) An isolated host cell which recombinantly expresses the haemopoietin receptor encoded by SEQ ID NO:3.
46. (Previously presented) The host cell of any one of claims 30, 36 or 45 wherein said host cell is an animal cell.
47. (Previously presented) A method of producing a recombinant polypeptide comprising culturing cells comprising the expression vector according to claim 10 for a time and under conditions sufficient to express a polypeptide encoded by the nucleic acid molecule as set forth in SEQ ID NO:3 in said expression vector and isolating said recombinant polypeptide.
48. (Currently amended) The isolated nucleic acid sequence of claim 1 ~~Claim 37~~ wherein said sequence consists ~~essentially~~ of nucleotides 142-1098 of SEQ ID NO: 3.
49. (Currently amended) The isolated nucleic acid sequence of claim 1 ~~Claim 37~~ wherein said sequence consists ~~essentially~~ of nucleotides 142-1338 of SEQ ID NO: 3.

50. (Currently amended) The isolated nucleic acid sequence of claim 1~~Claim 37~~ wherein said sequence consists ~~essentially~~ of nucleotides 142-414 of SEQ ID NO: 3.

51. (Currently amended) The isolated nucleic acid sequence of claim 1~~Claim 37~~ wherein said sequence consists ~~essentially~~ of nucleotides 415-1086 of SEQ ID NO: 3.

52. (Currently amended) The isolated nucleic acid molecule of claim 38 comprising a sequence of nucleotides encoding the amino acid sequence set forth in SEQ ID NO:4.